

Infection with the dinoflagellate parasite *Blastodinium* spp. in two Mediterranean copepods

Alf Skovgaard*

Departament de Biologia Marina i Oceanografia, Institut de Ciències del Mar, CMIMA, CSIC,
Passeig Marítim de la Barceloneta 37–49, 08003 Barcelona, Catalonia, Spain

ABSTRACT: Infection with the intestinal parasite *Blastodinium mangini* resulted in reduced survival of starved adults of the Mediterranean copepod *Oncaea* sp. No such effect was measurable for *Corycaeus* sp. infected by *Blastodinium navicula*. Both sexes of *Oncaea* sp. adults were infected by *B. mangini* and infected copepods were able to mate successfully. However, *Oncaea* sp. females infected by *B. mangini* did not produce eggs and, thus, appeared to be sterile. By inducing sterility, the harmful effect of *B. mangini* infection on *Oncaea* sp. populations is bound to be more profound than the effect on the individual level. The mechanism of infection by *Blastodinium* spp. remains unknown, but uptake of parasite zoospores by *Oncaea* sp. copepodites concurs with the hypothesis that infection occurs through the ingestion of zoospores by juvenile hosts.

KEY WORDS: Copepod · Parasite · Dinoflagellate · *Blastodinium* spp. · *Oncaea* sp. · *Corycaeus* sp.

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Copepods are a key link between microalgal primary production and grazing by larger pelagic organisms in the sea (e.g. Huys & Boxshall 1991). The marine planktonic food web is, however, not a single-strained food chain in which larger grazers simply feed on smaller organisms; some copepods are carnivores and feed on larger zooplankton organisms (Gophen & Harris 1984, Ohtsuka & Kubo 1991) and various microorganisms nourish themselves as parasites on their expected grazers such as copepods (Théodoridès 1989).

Dinoflagellates are particularly well represented among the parasites of copepods (Chatton 1920, Ho & Perkins 1985), but the consequences of parasitism for natural copepod communities have received little attention. Among the more common parasitic dinoflagellates of copepods in warmer seas are *Blastodinium* spp. A dozen species have been described that are all specialized inhabitants of the gut of marine planktonic copepods (Chatton 1920, Sewell 1951). The life cycle of *Blastodinium* spp. is only partially known. Briefly, a copepod is infested by a unicellular stage of the parasite, which then occupies the gut of the copepod, where

it grows and develops into a multicellular stage (the trophont) of considerable dimension (up to several 100 µm long, depending on the parasite species and size of the host organism). At maturity, numerous zoospores (dinospores) are released through the anus of the host. How the infection of a new host occurs is not known. It is believed that zoospores are ingested and then develop inside the gut of their host instead of being digested (Chatton 1920), but this mode of infection has never been confirmed.

Most *Blastodinium* species contain chloroplasts (Chatton 1920, Sewell 1951) and are able to perform photosynthesis (Pasternak et al. 1984), but whether the metabolic needs of the photosynthetic species are partly met by heterotrophy remains speculation. Limited information is available on the occurrence and infection frequencies of *Blastodinium* spp., since few investigations of copepod parasites have been done systematically. Most reports of *Blastodinium* spp. infections have been made in warm temperate or tropical waters where large proportions of copepods, for the most part adult females, have been found to contain these parasites (summarized by Shields 1994 and Coats 1999). Infection with *Blastodinium* spp. is

*Email: skovgaard@icm.csic.es

believed to have no severe physiological effect on the host, since most infected copepods observed have seemed in otherwise good condition, although infected adult females typically have reduced or disintegrated gonads and, consequently, are believed to be sterile (Chatton 1920, Sewell 1951). Based on this pathology, one may categorize *Blastodinium* spp. as a parasitic castrator rather than a true parasite (Kuris 1974).

The present study addresses the effect of infections by *Blastodinium mangini* and *B. navicula* on *Oncaea* sp. and *Corycaeus* sp., respectively (Copepoda, Cyclopoida). Species of these copepod genera frequently host *Blastodinium* spp. (Chatton 1920). While *B. mangini* predominantly infects *Oncaea* spp. (Chatton 1920, Sewell 1951), *Corycaeus giesbrechti* is the sole host of *B. navicula* (Chatton 1920, as *C. venustus*). The only exceptions to the latter host–parasite relationship are 2 specimens of *B. navicula* that, with some uncertainty, were found in *O. venusta* in the Arabian Sea (Sewell 1951).

According to Chatton (1920), morphological features of the trophont are important characters for identification of *Blastodinium* species. The trophont of *B. mangini* is 200 to 350 μm long with a cylindrical to fusiform shape. It is often vaguely curved and has rounded ends. The trophocyte is median to apical in position. The trophont of *B. navicula* is 150 to 200 μm long and is symmetrically fusiform with a distended central part and pointed ends. The trophocyte is submedian in position. Both species have brownish pigmentation and are usually gregarious.

MATERIALS AND METHODS

Sampling. Copepods were collected on August 19 (*Oncaea* sp.) and 25 (*Corycaeus* sp.), 2003, off Port Olímpic, Barcelona, Spain (sampling position: 41.22° 775' N, 02.13° 150' E; surface water temperature: 26°C). Samples were taken using a 100 μm mesh size plankton net pulled vertically from the bottom (38 m) to the surface. The live animals were kept at 26°C and all subsequent handling and experiments were also made at this temperature. On each sampling day, a parallel zooplankton sample was fixed immediately with 10% borax-buffered formalin and stored at 4°C in darkness. Infection frequencies of *Blastodinium* spp. in the fixed samples were determined by counting hosts with and without parasites using a stereomicroscope. All *Corycaeus* sp. in the sample were enumerated and dissected. Due to the high abundance of *Oncaea* sp., this sample was split into 10 equal aliquots of which only 1 was counted. *B. mangini* inside *Oncaea* sp. was visible without dissection of the animals.

Photographs of live copepods and formalin-fixed parasite trophonts were made using a Canon Power-

Shot G2 digital still camera mounted on an Olympus stereomicroscope or on a Zeiss Axiovert microscope with combined transmitted and epifluorescent illumination (filterset 09). Copepods of the genus *Oncaea* were identified using species descriptions by Heron & Bradford-Grieve (1995) and Böttger-Schnack (2001), and *Corycaeus* by use of the keys by Rose (1933) and Boxshall & Halsey (2004).

Survival of hosts. The survival times of starved adult copepods were determined to investigate whether *Blastodinium* spp.-infected individuals had different survival than healthy individuals, i.e. copepods without visible *Blastodinium* spp. infection. *Oncaea* sp. adults (148) were isolated, half of which were infected with *B. mangini*, and placed individually in 2 ml GF/F-filtered seawater in 3 ml Falcon™ multiwell cell culture plates (BD Biosciences). For each treatment (infected versus healthy), 50 copepods were placed in light (approx. 100 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) and 24 copepods were placed in darkness. For *Corycaeus* sp., 23 copepods were incubated for each treatment, all in the light. The number of live copepods was then monitored daily by observing each animal independently under a stereomicroscope. Immobile copepods were manipulated carefully with a fine needle to provoke mobility and thereby ascertain whether they were alive. Furthermore, the number of copepods that had produced eggs was noted, and also whether infected copepods had released *Blastodinium* spp. zoospores. Survival of infected and healthy copepods was analyzed using Kaplan-Meier plots and log-rank tests to compare the survival of 2 populations (SigmaStat® 3.0 software, Systat®). Dead copepods were fixed in formalin for species and sex determination.

Zoospore growth/survival. Experiments were also performed to determine whether *Blastodinium mangini* zoospores were able to grow or survive outside their hosts and whether *Oncaea* sp. grazed upon the zoospores. Zoospores for these experiments were obtained by incubating 70 to 80 *B. mangini*-infected *Oncaea* sp. for 24 h in a few ml of filtered seawater. After this incubation, the suspension of released zoospores was collected with a pipette. Aliquots containing 400 zoospores ml^{-1} were distributed into six 20 ml glass vials of which 3 were incubated in darkness and 3 were incubated in the light (approx. 100 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$). The concentration of zoospores was then monitored daily for 8 d by retrieving 1 ml from each vial and counting the number of cells after fixation in Lugol's solution, using an inverted microscope.

Grazing on zoospores. In order to see whether healthy *Oncaea* sp. grazed upon *Blastodinium mangini* zoospores, a suspension of zoospores in filtered seawater was incubated for 24 h in 5 different treatments that all consisted of 2 ml of zoospore sus-

pension ($300 \text{ cells ml}^{-1}$) contained in 3 ml wells (Falcon™ multiwell plates). The 5 treatments were: one with 2 adult *Oncaea* sp. females, one with 5 *Oncaea* sp. copepodites (cII–cIII, judged from their size), and one with 10 *Oncaea* sp. nauplii. The remaining 2 treatments consisted of one control without potential grazers and another with 2 *Acartia grani* females. The latter served as a 'positive control' to ensure that the experimental set-up was adequate to prove the occurrence of zoospore uptake. All treatments were set up in triplicate. *A. grani* originated from the culture collection at the Institut de Ciències del Mar, Barcelona, Spain. *Oncaea* sp. adults, copepodites and nauplii came from a temporary culture reared on a mixture of the dinoflagellate *Oxyrrhis marina* and the cryptophyte *Rhodomonas salina*. This culture, however, only survived for a few months, presumably because the food items were not adequate or because the copepods fed on their own offspring. Finally, a fraction of this temporary culture was transferred to a 600 ml culture tissue bottle, which then received a suspension of *B. mangini* zoospores within a day after these had been released from a dozen freshly collected infected *Oncaea* sp. In order to see whether the addition of zoospores would lead to visible infection in previously uninfected hosts, this subculture was checked daily

by observing the copepods inside the culture tissue bottle using a stereomicroscope equipped with dark-field illumination.

RESULTS

Identification of *Blastodinium* spp.

When viewed with a stereomicroscope with transmitted bright-field illumination, infection of *Blastodinium mangini* in live *Oncaea* sp. was relatively easily recognized as one or more dark brownish-greenish bodies, trophonts, in the copepod's gut (Fig. 1A). Recognition was facilitated by use of dark-field illumination by which the parasite trophonts gained a distinctive yellowish colouration (Fig. 1B). Dissection of infected *Oncaea* sp. revealed the typical, slightly fusiform, shape of the *B. mangini* trophont (Fig. 1C). The ends of the trophonts were rounded and the trophocytes were median to apical in position (apical in Fig. 1C). Each infected host usually had 2 to 4 trophonts. Due to the chloroplasts they contained, the parasites exhibited red autofluorescence when viewed under epifluorescence microscopy, which assisted in detecting and confirming *B. mangini* infection without




Fig. 1. (A,B) *Blastodinium mangini* in *Oncaea* sp., live organisms. (A) Bright-field image of infected female. (B) Dark-field image of infected female; 2 to 3 *B. mangini* trophonts are visible as yellowish rods inside the copepod's gut. (C,D) Formalin-fixed *Blastodinium* spp. trophonts. Colour has disappeared due to fixation. (C) *B. mangini* isolated from its host, *Oncaea* sp. (D) *B. navicula* isolated from *Corycaeus giesbrechti*. Arrowheads point to the extremes of trophocytes. (A) and (B) are of equal magnification, and (C) and (D) are of equal magnification

dissection of the hosts (Fig. 2). The characteristic colours and fluorescence of *Blastodinium* spp. was, however, only easily visible in fresh (live) hosts or in formalin-fixed hosts that had been stored cold (at 4°C) in darkness. In formalin-fixed hosts that had been stored at room temperature and exposed to daylight for

longer periods, parasites were difficult to discern due to poor preservation of chlorophyll, and the contour of the parasites was the only characteristic whereby they could be distinguished. Total infection frequency of *Oncaea* sp. in the parallel formalin-fixed sample was 2.0% (792 copepods). For adult females, males and




Fig. 2. *Oncaea* sp. infected with *Blastodinium mangini*, live organisms. (A) Non-infected female. (B) Infected female, *B. mangini* trophonts visible as dark rods in the gut of the host. (C) Same specimen as (B), but viewed with epifluorescence microscopy illustrating chlorophyll content of 4 parasite trophonts. (D) Female *B. mangini*-infected *Oncaea* sp. carrying eggs. Epifluorescence microscopy. (E) Copulating *Oncaea* sp. couple in which both parts are infected with *B. mangini*. Epifluorescence microscopy. (F) Infected *Oncaea* sp. female with attached spermatophores. Epifluorescence microscopy. (A) to (D) are of equal magnification, and (E) and (F) are of equal magnification

late-stage copepodites the infection frequencies were 3.7, 1.5 and 1.1%, respectively.

The *Corycaeus* sp. specimens from the experiment, which were examined for species determination, were all *C. giesbrechti*. All *Blastodinium* sp.-infected *Corycaeus* sp. recognized in the formalin-fixed sample were, likewise, *C. giesbrechti*, and in all cases the parasite had the size and shape of the trophont characteristic for *B. navicula*, i.e. the shape was symmetric with pointed ends and the trophocyte was close to median in position (Fig. 1D). There was always more than one trophont in each infected host. Five out of 51 adult *Corycaeus* sp. in the quantitative, formalin-fixed sample were infected by *B. navicula*. Due to the opaqueness of the fixed host, these parasites were barely recognizable without dissecting the copepod even when the sample had been stored cold and dark.

Survival of hosts

Oncaea sp. infected with *Blastodinium mangini* lived significantly shorter than healthy *Oncaea* sp. (Fig. 3, Table 1). Healthy, starved *Oncaea* sp. stayed alive on average 10.7 d, whereas *B. mangini*-infected individuals survived on average only 7.4 d. Median survival time for healthy and infected *Oncaea* sp. was 10 and 7 d, respectively. When comparing healthy and *B. mangini*-infected copepods separately, there was no difference in survival times with respect to whether

Table 1. Survival of starved *Oncaea* sp. and *Corycaeus* sp. infected with *Blastodinium mangini* and *B. navicula*, respectively, compared with healthy (non-infected) copepods. Different superscript letter denotes that survival curves of populations are statistically different (Kaplan-Meier plots; log rank test, $p < 0.001$)

Treatment	<i>Oncaea</i> sp.		<i>Corycaeus</i> sp.	
	Survival time (d)	SE	Survival time (d)	SE
Total				
Infected	7.4 ^a	0.2	2.8 ^c	0.2
Non-infected	10.5 ^b	0.4	3.3 ^c	0.3
Infected				
Light	7.2 ^a	0.4	–	–
Dark	8.1 ^a	0.4	–	–
Non-infected				
Light	10.3 ^b	0.5	–	–
Dark	11.2 ^b	1.3	–	–

the animals were kept in light or in darkness (Table 1, survival curves not shown).

Mean survival time for healthy and *Blastodinium navicula*-infected *Corycaeus* sp. was 3.3 and 2.8 d, respectively (Fig. 4, Table 1) and was, thus, considerably shorter than that of *Oncaea* sp. There was no significant difference between survival times of healthy versus *B. navicula*-infected individuals of *Corycaeus* sp. (Fig. 4, Table 1). Median survival time was 4 d for healthy individuals and 3 d for infected individuals.




Fig. 3. Kaplan-Meier survival curves of *Blastodinium mangini*-infected (●) and healthy (non-infected, ○) *Oncaea* sp. Initial number of copepods was 74 from each treatment. Error bars represent ± SE




Fig. 4. Kaplan-Meier survival curves of *Blastodinium navicula*-infected (●) and healthy (non-infected, ○) *Corycaeus* sp. Initial number of copepods was 23 from each treatment. Error bars represent ± SE

Egg production

The majority of both healthy and infected *Oncaea* sp. used for the survival experiment were adult females (78 and 54 %, respectively), but they did not carry egg sacs when the experiment was initiated. Of the healthy females, 55 % produced eggs within the first 5 d of the experiment (Fig. 5), whereas none of the infected females produced eggs during the experiment. All but 2 infected *Oncaea* sp. released zoospores within the first 3 d of the survival experiment (Fig. 5). The remaining 2 infected copepods never released zoospores, but epifluorescence microscopy confirmed that they were infected (as in Fig. 2C–F). The starved infected animals were, thus, able to stay alive for up to 1 wk after the parasite had completely or partially left the host. On a single occasion a *Blastodinium mangini*-infected *Oncaea* sp. female from a field sample was observed to carry eggs (Fig. 2D), but the eggs never hatched. Male and female *B. mangini*-infected *Oncaea* sp. that had been placed together in filtered seawater in a Petri dish were seen mating on several occasions (Fig. 2E). Some of these females, as well as several infected females collected in the field, had attached spermatophores (Fig. 2F).

Zoospore growth/survival

Free-swimming *Blastodinium mangini* zoospores originating from *Oncaea* sp. did not exhibit photosynthetic growth when incubated in *f/2*-enriched sea-




Fig. 5. Cumulative number of *Blastodinium mangini*-infected *Oncaea* sp. that had produced zoospores and healthy (non-infected) *Oncaea* sp. that had produced eggs during the survival experiment shown in Fig. 3

water. Cell numbers declined during the incubation period (Fig. 6), and this decrease was not evidently dependent on whether cells were incubated in light or in darkness.

Grazing on zoospores

No feeding on zoospores by *Oncaea* sp. adults or nauplii was detectable, but during the incubation there was a reduction in the number of free-swimming *Blastodinium mangini* zoospores in the treatments with *Oncaea* sp. copepodites and *Acartia grani* (Fig. 7). However, the attempt to induce infection by adding freshly released *B. mangini* zoospores to a temporary *Oncaea* sp. culture was unsuccessful, since no signs of re-infection were detectable for the time period the *Oncaea* sp. culture was observed (4 wk).

DISCUSSION

Taxonomy of parasites and hosts

All *Oncaea* sp. used in the experiments had comparable gross morphology. Identification of the specimens was hampered due to their condition (they were dead at the time of fixation). However, the females that were successfully dissected were clearly members of the *O. media* species complex. They were smaller (body length < 0.6 mm) than *O. media* and the propor-




Fig. 6. Development of numbers of *Blastodinium mangini* zoospores (previously released from *Oncaea* sp.) incubated in *f/2*-medium in the light ($100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) and in the dark. Error bars represent \pm SE




Fig. 7. Potential grazing on *Blastodinium mangini* zoospores by *Oncaea* sp. adults, nauplii and copepodites, and by *Acartia grani*. Error bars represent +SE. *: concentration of zoospores after 24 h was significantly different from the control without grazers (t -test, $p < 0.01$)

tional lengths of their urosome segments indicated a close similarity to the 2 sibling species *O. scottodiarloi* and *O. waldemari*. Other more distinct species of *Oncaea* were present in low numbers in the collected samples. These were distinguishable due to differences in size and gross morphology, and were not used for the experiments. Dissection of infected specimens (Fig. 1C) confirmed that the parasite of *Oncaea* sp. was *Blastodinium mangini*. The size and shape of the trophont as well as the trophocyte location were typical for this species (Chatton 1920). A variety, *B. mangini* var. *oncaea*, has been described from *O. media* (Chatton 1920). However, it is not always possible to distinguish the variety from the typical form (Sewell 1951) and in the present study no attempt was made to identify *B. mangini* further than to the species level. Neither *O. scottodiarloi* nor *O. waldemari* have been reported as hosts for *B. mangini* before. However, both species have only recently been recognized as separate species (Bersano & Boxshall 1994, Heron & Bradford-Grieve 1995) and they may in the past have been reported as the related species *O. media* (Böttger-Schnack 2001). Further studies are needed to determine exactly which *Oncaea* species are hosts for *Blastodinium* spp., incorporating recent knowledge on *Oncaea* species diversity.

There might have been more than one *Corycaeus* species in this study since not all animals were dissected and examined for species determination, and the subgenus *Onychocorycaeus*, to which *C. giesbrechti* belongs, contains species with gross morphologies comparable to *C. giesbrechti*. At the end of the survival experiment, the state of the parasite of *Corycaeus* did not allow confirmation of the parasite species due to sporulation and decomposition of the parasite trophont. Species determina-

tion of *Blastodinium* inside a live *Corycaeus* sp. is not feasible due to the limited transparency of this host species. However, infected *Corycaeus* sp. specimens that were taken from the parallel, formalin-fixed sample were dissected and all contained the parasite *B. navicula*. The size and symmetrically pointed shape of the trophont and the submedian location of its trophocyte (Fig. 2D) are all typical characteristics for this species (Chatton 1920). The host was in all cases *C. giesbrechti*, which is the only member of the subgenus *Onychocorycaeus* that has been found to host *Blastodinium* spp. Furthermore, *B. navicula* is the only *Blastodinium* species known to infect this subgenus in the Mediterranean Sea (Chatton 1920), and it is therefore plausible that all the infected *Corycaeus* spp. in this study were *C. giesbrechti* and that the parasite was in all cases *B. navicula*.

Effects of infection

The survival experiments validate previous assumptions that infection by *Blastodinium* spp. is not directly lethal to its hosts (Chatton 1920, Sewell 1951). However, infection by *B. mangini* in *Oncaea* sp. leads to decreased survival of the host (Fig. 3). This negative effect on survival may be a direct effect on the host's physiology, but it is also possible that infection with *Blastodinium* spp. reduces the grazing capability of the host or that the parasite exploits the gut content of its host. In the latter 2 cases the negative effect on the host would be indirect by reducing the nutritional uptake of the host. In this respect it would be interesting to study the feeding biology of *Blastodinium* spp.-infected copepods. No effect of *B. navicula* infection was evident for *Corycaeus* sp. Since starved *Corycaeus* sp. only survived for a few days, it is possible that a larger number of animals would be needed to show any effect by the experimental set-up used here. A study on the effect of *Blastodinium* sp. infection in *Eucalanus subtenius* yielded ambiguous results on host survival (Pasternak et al. 1984). However, some copepods in that study became 'heavily overgrown with infusorians' and this might have affected their viability.

One may hypothesize that infection with *Blastodinium* spp. could have a favourable effect on its host by providing it with photosynthate (Pasternak et al. 1984). This hypothesis is, however, not supported by the results listed in Table 1: starved, infected *Oncaea* sp. did not benefit from being incubated in the light, where one would have expected that hosting a photosynthetic symbiont would have been an obvious advantage. A similar conclusion may be drawn from the data given by Pasternak et al. (1984), who concluded that the amount of organic matter *Blastodinium* spp. can deliver to its host is negligible.

Host sterility

Previous studies on the fecundity of *Blastodinium* spp.-infected copepods have been somewhat conflicting. It is generally believed that infected copepod females are sterile or at least have reduced fecundity since they have been observed to have poorly developed or disintegrated ovaries (Chatton 1920, Sewell 1951). However, examples also exist where *Blastodinium* spp.-infected females have seemed to have normally developed gonads (Ianora et al. 1990). Sterility of *Blastodinium* spp.-infected females is supported by experimental data in the present study, in which more than half of the incubated healthy females produced eggs within the first 5 d of incubation (Fig. 5), whereas none of the infected females produced eggs. It is unknown how many of the incubated females were fertilized prior to the experiment, but infected and healthy females may have been fertilized at equal ratios, since infected females are also able to mate (Chatton 1920, Sewell 1951; Fig. 2E,F). Hence, *Blastodinium* spp.-infected females may mate successfully, but mating does not lead to egg production, probably because the gonads are poorly developed. If infected females do not feed, the lack of egg production may be because the gonads of these females stay immature. This has been shown for a starved calanoid copepod (Niehoff 2000). On rare occasions, *Oncaea* sp. females infected with *B. mangini* do produce eggs (Fig. 2D). It is possible that the fertility of an infected female is dependent on the developmental stage of the female when it is infected. A female infected at a late developmental stage may, thus, have developed ovaries (Ianora et al. 1990), but a copepod female infected with *Blastodinium* spp. has never been observed to produce offspring and may, nevertheless, be sterile. Whether a copepod is able to fully recover an infection with *Blastodinium* spp. and regain its fecundity is a possibility that needs to be investigated. Adult male copepods are generally not infected by *Blastodinium* spp. (Chatton 1920, Sewell 1951). *Oncaea* is an exception to this rule, since both adult males and females of this genus are infected by *B. mangini* (Sewell 1951; Fig. 2E). Whether infected males are fertile is unknown.

Infection via ingestion

Attempts to induce an infection of dinoflagellate parasites in copepods under laboratory conditions have so far been unsuccessful (Kimmerer & McKinnon 1990, the present study), and how new hosts are infected remains an enigma. The most plausible

hypothesis is that a parasite zoospore is ingested by a copepod and, in the case of *Blastodinium* spp., develops inside the hosts' gut. A prerequisite for this would be that zoospores are actually ingested by their future host. The feeding biology of *Oncaea* is not well understood, but some species of the genus are known to be raptorial predators on other zooplankton organisms, in particular appendicularians and their houses (Ohtsuka et al. 1996, Go et al. 1998). Hence, it is unlikely that adults of *Oncaea* sp. would ingest prey as small as a *Blastodinium mangini* zoospore (equivalent spherical diameter ~10 µm) and in fact this was not the case in the feeding experiment (Fig. 7). However, the smaller copepodites of *Oncaea* sp. reduced the number of free-swimming zoospores, and it is possible that parasitism by *B. mangini* is established in juvenile hosts that, probably, depend on food items different from the diet of adults. This would confirm a hypothesis put forward by Chatton (1920) that *Blastodinium* might infect nauplii and/or early copepodite stages rather than adult copepods. It cannot be excluded, however, that zoospores may enter a new host actively, e.g. through the anus, or that they need to go into a yet unknown life cycle stage (e.g. cysts) to be able to establish a new infection. The possibility also exists that the parasite is transmitted to adult copepods through an intermediate host, but this possibility has never been explored.

CONCLUSION

Infection with *Blastodinium* spp. is less drastic for the individual copepod than, e.g., infection with *Syn- dinium*, a parasitic dinoflagellate genus that kills its host by literally devouring it from inside (Chatton 1910, Kimmerer & McKinnon 1990). However, copepods infected with *Blastodinium* spp. do suffer from reduced fitness and may also have a shorter lifespan in nature. In addition, parasite-induced sterility in females may have considerable effects on the copepod population not only because infected females do not produce eggs, but also because mating with a sterile female is bound to reduce the overall potential for successful mating of healthy males. Sterility of a host may thus have a more important effect on the host population than does the death of the infected host (Kuris 1974). The possible impact of sterile mating in copepods remains to be investigated.

Acknowledgements. I thank E. Saiz for constructive comments and discussion and R. Böttger-Schnack for advice on *Oncaea* taxonomy. This work was financed by the European Commission, Directorate General Research, through a Marie Curie Postdoctoral fellowship (contract no. HPMF-CT-2002-01931).

LITERATURE CITED

- Bersano JGF, Boxshall GA (1994) Planktonic copepods of the genus *Oncaea* Phillipi (Poecilostomatoida: Oncaeidae) from the waters off southern Brazil. *Nauplius*, Rio Grande 2:29–41
- Böttger-Schnack R (2001) Taxonomy of Oncaeidae (Copepoda, Poecilostomatoida) from the Red Sea. II. Seven species of *Oncaea* s.str. *Bull Nat Hist Mus Lond (Zool)* 67: 25–84
- Boxshall GA, Halsey SH (2004) An introduction to copepod diversity. The Ray Society, London
- Chatton É (1910) Sur l'existence de dinoflagellés cœlomiques. Les *Syndinium* chez les copépods pélagiques. *C R Hebd Séanc Acad Sci Paris* 102:645–656
- Chatton É (1920) Les Péridiniens parasites. Morphologie, reproduction, éthologie. *Arch Zool Exp Gén* 59:1–475, plates I–XVIII
- Coats DW (1999) Parasitic life styles of marine dinoflagellates. *J Eukaryot Microbiol* 46:402–409
- Go YB, Oh BC, Terazaki M (1998) Feeding behavior of the poecilostomatoid copepods *Oncaea* spp. on chaetognaths. *J Mar Syst* 15:475–482
- Gophen M, Harris RP (1984) Visual predation by a marine cyclopoid copepod, *Corycaeus anglicus*. *J Mar Biol Assoc UK* 61:391–399
- Heron GA, Bradford-Grieve J (1995) The marine fauna of New Zealand: pelagic Copepoda: Poecilostomatoida: Oncaeidae. *N Z Oceanogr Inst Mem* 104:1–57
- Ho J, Perkins PS (1985) Symbionts of marine copepoda: an overview. *Bull Mar Sci* 37:586–598
- Huys R, Boxshall GA (1991) Copepod evolution. The Ray Society, London
- Ianora A, Scotto di Carlo B, Mazzocchi MG, Mascellaro P (1990) Histomorphological changes in the reproductive condition of parasitized marine planktonic copepods. *J Plankton Res* 12:249–258
- Kimmerer WJ, McKinnon AD (1990) High mortality in a copepod population caused by a parasitic dinoflagellate. *Mar Biol* 107:449–452
- Kuris AM (1974) Trophic interactions: similarity of parasitic castrators to parasitoids. *Q Rev Biol* 49:129–148
- Niehoff B (2000) Effect of starvation on the reproductive potential of *Calanus finmarchicus*. *ICES J Mar Sci* 57: 1764–1772
- Ohtsuka S, Kubo N (1991) Larvaceans and their houses as important food for some pelagic copepods. *Proc 4th Int Conf Copepoda*, Karuizawa, Japan, 16–20 September 1990. *Bull Plankton Soc Jpn Spec Vol*:535–551
- Ohtsuka S, Böttger-Schnack R, Okada M, Onbé T (1996) In situ feeding habits of *Oncaea* (Copepoda: Poecilostomatoida) from the upper 250 m of the central Red Sea, with special reference to consumption of appendicularians houses. *Bull Plankton Soc Jpn* 43:89–105
- Pasternak AF, Arashkevich YG, Sorokin YS (1984) The role of the parasitic algal genus *Blastodinium* in the ecology of planktonic copepods. *Oceanology* 24:748–751
- Rose M (1933) Copépodes pélagiques. In: *Faune de France*. Lechevalier, Paris
- Sewell RBS (1951) The epibionts and parasites of the planktonic Copepoda of the Arabian Sea. *John Murray Expedition 1933–34. Sci Rep Br Mus Nat Hist* 9:255–394
- Shields JD (1994) The parasitic dinoflagellates of marine crustaceans. *Annu Rev Fish Dis* 4:241–271
- Théodoridès J (1989) Parasitology of marine zooplankton. *Adv Mar Biol* 25:117–177

*Editorial responsibility: John Dolan,
Villefranche-sur-Mer, France*

*Submitted: July 16, 2004; Accepted: November 15, 2004
Proofs received from author(s): January 12, 2005*